

# INTEGRABILITY PROBLEM

$$f: M \rightarrow M \quad p.h \quad TM = \underbrace{E^3 \oplus E^C \oplus E^u}_{E^{cs} \oplus E^{cu}}$$

$f$  is dynamically coherent

$\exists f$ -inv.  $W^{cs}$  &  $W^{cu}$  tangent

to  $E^{cs}$  and  $E^{cu}$ .

$E^s$  and  $E^u$  are uniquely integrable

Prop (Buraqo-Ivanov)  $\gamma$  curve  
tg to  $E^{cs}$  and transverse to  $E^s$   
then  $S = \cup W^s(\gamma)$  is a surface  
tg to  $E^{cs}$  yes

Idea:  $C^0$ -frobienius property.

Thm (Buražo-Ivanov)

Assume  $E^0$  are orientable, DF-pres.

$\exists f^{-cs}, f^{cu}$   $f$ -inv branching


foliations and tangent to

$E^{cs}$  and  $E^{cu}$ .

Idea: Choose the "lowest"  
surfaces

→  $f$ -inv.

→ no "topological crossings"



# Branching Foliation:

$\mathcal{F}$  collection  $\{L_\alpha\}$  of complete surfaces tangent to a continuous dist.

$E$  s.t:  $\rightarrow \cup L_\alpha = M$

$\rightarrow$  no top. crossings.

$\rightarrow X_n \rightarrow X, X_n \in L_{\alpha_n}$


$L_{\alpha_n} \rightarrow L_\alpha$

Remark/Deg (Bonatti-Wilkinson)

A foliation is a branching

foliation without branching.

~~It~~

If  $\exists M^c$  then  $S^3$  cannot  
admit  $\overline{p.h.}$  diffeos. 

[Thm (BI) It is possible to  
"blow up" the branching foliation]

Corollary  $S^3$  does not admit p.h. diffeos

# Naive Idea:



Under 2 assumptions this works:

- Brin
- Absolute partial hyperbolicity
  - quasi-isometry of  $\mathcal{N}^u$ .



[ Thm (BBT)  $f: \mathbb{T}^3 \rightarrow \mathbb{T}^3$  absolute  
partially hyp. diffeo  $\Rightarrow f$  is  
dynamically coherent

Proof: Showing  $\mathcal{W}^s, \mathcal{W}^u$  are  
quasi-isometric.  
 $d_{\tilde{M}}(x, y) \leq a d(x, y) + b$

Thm (J, F. Rodrigues Horta, Ures)  
∃ open sets of p.h. in  $\mathbb{T}^3$   
which are not d.c.

Conj (RH, RH, U) IF  $\nexists T$   $\delta$ -periodic  
tangent to  $E^s$  or  $E^{cu} \Rightarrow f$  is d.c.


# Example:

$A: \mathbb{T}^2 \rightarrow \mathbb{T}^2$  linear diffeo

eigenvalues  $\lambda < 1$  and  $\lambda^{-1}$

$f_1: \mathbb{T}^2 \times [1, 17] \rightarrow \mathbb{T}^2$   $f_1(x, t) = (Ax, \mu t)$


$\mu < 1$




$$f_2: \mathbb{T}^2 \times [-1, 1] \rightarrow \mathbb{S}^1 \quad f_2(x, t) = (Ax, \theta t)$$


---

$$f: \mathbb{T}^2 \times \mathbb{S}^1 \rightarrow \mathbb{S}^1 \quad f(x, t) = (Ax + \psi(t), v^s)$$



$\psi(t)$





TCSA

TCSA

The example can be done

$$\text{in } M_A = \mathbb{T}^2 \times [0, 1] / \sim$$

[Thm (HP) If  $\beta T$  is on cu  
 $\Rightarrow f$  is dyn. coherent.  
(if  $\pi(M)$  is solvable)

$\rightarrow$  Transitivity, conservative  
 $\rightarrow$  absolutely p.h.

Prop If  $\exists$  periodic cs on  $\alpha$   
 $\neg T \Rightarrow f$  is not abs. p.h.





1)  $f/T$  where  $T$  is CS.

$$\lambda_{\text{top}}(f/T) = \lambda_{\text{top}}(A)$$

var. ppe

Ruelle meq }  $\Rightarrow \exists \mu$

$$\lambda^c(\mu) > \lambda_{\text{top}}(A) - \varepsilon$$

2)  $M^u$ -saturated

~~-----~~

$\Lambda$  p.h. attractor

~~$\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}$~~   $C^s$   $f/\Lambda$  semiconj to  $A$

$$h_{\text{top}}(f/\Lambda) = h_{\text{top}}(A)$$

$\exists \mu'$  such that  $\lambda^u(x^i) \leq h_{\text{top}}(A)$